



I'm not robot



Continue

## Beplex forte drug information

Drugs & Drugs Used Side Effects Precautions Interaction Overdose Pictures Uses Side Effects Precautionary Measures Interactions Overdose Pictures Use This drug is a multivitamin product that is used to treat or prevent vitamin deficiency due to poor nutrition, certain diseases, or during pregnancy. Vitamins are an important building block of the body and help you maintain good health. Take this medicine for your mouth, usually once a day or as directed. Follow all instructions for packing the product or follow your doctor's instructions. Do not take more than the recommended dose. If you have any questions, ask your doctor or pharmacist. Take these medications regularly to get the most benefit from it. To help you remember, take it at the same time each day. Side effects Constipation, diarrhea or upset stomach may occur. These effects are usually temporary and may disappear when your body adjusts to this medicine. If any of these effects persist or get worse, talk to your doctor or pharmacist immediately. If you have been prescribed this medicine, remember that the benefit to you is greater than the risk of side effects. Many people who use these drugs do not have serious side effects. A very serious allergic reaction to this drug is rare. However, seek immediate medical advice if you notice any of the following symptoms of a serious allergic reaction: rash, itching/swelling (especially face/tongue/throat), severe dizziness, difficulty breathing. This is not a complete list of possible side effects. If you notice any other effects not listed above, talk to your doctor or pharmacist. In the U.S.- Call your doctor for medical advice on side effects. You can report side effects at the FDA on 1-800-FDA-1088 or in www.fda.gov/medwatch. In Canada – Call your doctor for medical advice on side effects. You can report side effects at Health Canada at 1-866-234-2345. Precautions Please inform your doctor or pharmacist before taking this medicine if you are allergic to any of the other ingredients: or soy beetle/peanut found in certain brands; or if you have other allergies. This product may contain inactive ingredients that may cause allergic reactions or other problems. Ask your pharmacist for more details. Before taking this medicine, tell your doctor or pharmacist about your medical history, in particular: alcohol use/abuse, liver problems, stomach/bowel problems (e.g. ulcer, colitis). If your brand name multivitamin also contains folic acid, talk to your doctor or pharmacist before taking it if you have vitamin B12 deficiency (pernicious anemia). Folic acid can affect some laboratory tests for vitamin B12 deficiency without treating this anemia. The lack of untreated vitamin B12 can cause serious problems with (e.g. peripheral neuropathy). Talk to your doctor or pharmacist for details. These medicines are treated in breast milk. Talk to your doctor before breast-feeding. Doctor. Interactions with medicines may alter the way your medicines work or increase the risk of serious side effects. This document does not contain all possible drug interactions. Keep a list of all products you use (including prescription/non-prescription medicines and herbal products) and share it with your doctor and pharmacist. Do not start, stop or change the dose of any medicines without a doctor. Some products that can interact with this drug are: other vitamin/dietary supplements. If your brand name multivitamin also contains iron, avoid taking this product at the same time as antacids, bisphosphonates (such as alendronate), levodopa, thyroid medicines (such as levothyroxine) or some antibiotics (such as tetracyclines such as cyproflaxacin). Ask your doctor or pharmacist how long you should wait between doses and to help find a dosage schedule that will work with all medications. If your brand name multivitamin also contains folic acid, be sure to tell your doctor or pharmacist if you are taking certain anti-seizure medicines (including hidantoins such as phenytoin). These drugs can interfere with some laboratory tests, can cause false test results. Make sure that the laboratory staff and all your doctors know that you are using this medicine. Overdose If someone has overdose and has serious symptoms such as fainting or difficulty breathing, call 911. Otherwise, call the venom control center immediately. U.S. residents can call their local poison control center at 1-800-222-1222. The people of Canada can call it a provincial poison control center. Symptoms of overdose may include: stomach pain, nausea, vomiting, diarrhea. Comments If your doctor has prescribed this medicine, do not share it. Check doctors and laboratories regularly. Some brands may also contain ingredients such as docusate. If you have questions about the ingredients in the brand name, ask your doctor or pharmacist. This product is not a substitute for proper nutrition. Remember that it is best to get your vitamins from healthy foods. Maintain a well-balanced diet and follow all dietary guidelines as directed by your doctor. Missed a dose If you are taking the medicine according to the prescribed schedule and miss a dose, take it as soon as you remember. If it is at the time of the next dose, skip the dose. Take your next dose at regular times. Do not double the dose to catch up. Storage/Store at room temperature between 59-86 degrees F (15-30 degrees C) away from light and moisture. Do not store in the bathroom. Not all medicines should be used for children and pets. Do not rinse the medicine in the toilet or put it in the drain unless you are told to do so. Dispose of this product correctly when it has expired or you no longer need it. For more details on how to dispose of the product safely, contact your pharmacist or pharmacist information last amended in May 2020. Copyright(c) 2020 First Databank, Inc. Images Sorry, No pictures are available for these medicines. Images. Resources for Beplex Tablet Find lowest prices on Verified Interactions Read Reviews (1) Selected between data included with permission and copyright protected by First Databank, Inc. This copyrighted material has been transferred from the licensed data provider and is not intended for distribution, unless permitted by applicable terms of use. TERMS OF USE: The information in this database is intended to complement, not replace, the expertise and judgement of health professionals. This information is not intended to cover all possible uses, instructions, precautions, drug interactions or side effects, nor should it be stated that the use of a particular medicinal product is safe, appropriate or effective for you or anyone else. Before taking any medicine, a healthcare professional should be consulted, any diet should be changed or any course of treatment started or discontinued. Beplex forte is used primarily to prevent or treat conditions where there is an increased requirement of vitamins and minerals, as in cases of anemia. Hearth and hepatic complications are the main side effects with high repeated doses of beplex forte. In the case of patients with hepatic impairment and cases of ulceration in optics should be completely avoided. Composition of Beplex Forte – Vitamin B9 (Folic acid) 1.5 MG, Vitamin B12 (cyanocobalamin) 15 MCG, Vitamin B3 (Niacinamide) 75 MG, calcium pantothenate 50 MG, Vitamin B6 (pyridoxine) 3 MG, Vitamin C (Ascorbic acid) 150 MG, Vitamin B2 (Riboflavin) 10 MG, Vitamin B1 (Thiamin) 10 MG, Vitamin B7 (Biotin) 260 MCG, 50 MG, Elemental Magnesium 32.4 MGManufactured by – Anglo-French Drugs and Industries Ltd.Recipe – Not Required (Available as OTC)Form – Tablets, Sirup and Injections.Price – Rs 29.22 for 20 Tablets.Expiry/Shell Life – 24 months from the date of manufacturingType of the drug – Multivitamin i Multi-Minerals Searchtano on hindi o Beplex Forte Using Beplex Forte Beplex Forte serves great goal in preventing and tipening various conditions It is prescribed for the following : Ulcers and innate language : It is used to treat oral ulcer and inflamed tongue due to vitamin deficiency. Hair problems: It is used to treat hair and grey hair fall by providing adequate amounts of hair follicle care and also treatment of weak nails. Anemia: Used to treat various types of anemia, such as pernicious anemia (vitamin B12 deficiency), Megaloblastic anemia (folic acid deficiency) etc. Vitamin and mineral deficiency: It is used to treat diseases resulting from vitamin and mineral deficiency such as Scurvy (vitamin C deficiency) and calcium deficiency and promote wound healing by providing adequate amounts of calcium. Nerve disorders: It is used to treat nerve diseases such as neuralgia (pain due to nerve damage) and Paraesthesia (a changed taste sensation). Poor dietary intake: Used to treat poor intake in cases of chronic alcoholics or elderly patients. Post-surgical cases: Used in patients with post-surgical problems. It is used to treat high cholesterol levels. Other: Used in cases to treat driers, muscle spasms and chronic gastrointestinal disorders. Read also: Betadin Used | Betahistine Uses How Does Beplex Forte Work? Beplex forte works by providing the body with essential nutrients in an appropriate quantity. Vitamin B complex and vitamin C deficiency can lead to a variety of diseases, such as delayed wound healing and make the body more lined up infections. It works by providing the body with vitamin C, which acts as an antioxidant, booster immunodeficiency system that can help reduce the risk of chronic diseases such as heart disease and cancer, allows the body to keep white blood cells to fight infections and is necessary for synthesis and use of certain amino acids and collagen, which helps to heal ranaBiotin Beplex forte is a vitamin soluble in water that is part of the vitamin B family. It helps the body convert certain nutrients into energy and also plays an important role in the health of hair, skin and nails. Vitamin B12 helps keep nerves and blood cells healthy and helps to synthesise DNA in all cells. It also helps to prevent megaloblastic anemia, which makes people tired and weak. How to take Beplex Forte? Beplex forte is available as tablets mainly Take the tablet peroralo with or without water, preferably with a meal, as it helps with rapid absorption. Do not chew or crush the tablets, but swallow the tablets as a whole. Do not change the dose in your own convenience. It is recommended that the patient goes through the instructions inside the package thoroughly in order to better understand the drug. Common Dose for Beplex Forte Dose should be decided by a doctor for each individual, such as age, weight, mental state, allergic history of the patient. Dose for adults: One tablet daily or twice a day in some cases, as advised by a doctor according to the severity of the conditionOdithage of the child: Not recommended for children, Pediatric consultation is recommended. The product should be taken at the same intervals so that an adequate amount of the medicine is present in the body at all times. Q. What happens if you overdose beplex forte? Beplex Forte should always be taken after the prescribed dose. Overdose may cause some serious undesirable effects. In the event of an overdose, a doctor or qualified healthcare professional should be consulted immediately. Beplex Forte should always be taken according to a certain timetable and, in case of a dose, try to take it as soon as you remember. However, avoid taking a missed dose if it is time to take a second dose as it may lead to drug toxicity or drug overdose. Q. What happens if i sing an expired Beplex Forte? Beplex Forte that has expired should not cause side effects, but it is recommended to avoid taking expired as it may not be effective enough to ensure that the should be immediately consulted in case of adverse symptoms after ingestion of the product that has expired. Q. What is the time to set up Beplex Forte? The patient may start to feel recovering within the first few days after treatment with Beplex forte. The dose and duration of the drug, as advised by the doctor, should be strictly observed for the complete elimination of all symptoms of Q. Until when do the effects of Beplex Forte remain? The time taken by the drug to show its effect may vary with each individual. Read also: Dose of azithromycin | Bekosul Capsule Dosage | Banocide Forte Dosage When to Avoid Beplex Forte? Do not take Beplex Forte tablets in the following scenarios Liver disease: In known cases of moderate to severe liver dysfunction or disease, beplex forte should only be avoided or used if your doctor tells you to. Allergy: In known cases of allergy to Beplex Forte or any of the ingredients of the drug. Breastfeeding women: In the case of a breastfeeding mother or breastfeeding mother, Beplex Forte is avoided. Peptic ulcer: Patients with peptic ulcer should be avoided. Cardiac arrhythmias: Patients with cardiac arrhythmia should be avoided. Precautions While taking Beplex Forte Long-term use: Do not use it for long periods of time unless your doctor has told you to. Allergic reaction: An allergic reaction if it occurs after taking Beplex Forte, the doctor or qualified medical staff should be informed immediately. Avoid changing the dose: changes to the medicine should be avoided unless your doctor has told you. Concurrent products: Do not mix it with other dietary supplements. Warnings Beplex Forte Beplex forte, which is prescribed with antibiotics, is used to deter side effects such as oral ulcers, diarrhoea, etc. Do not change the dose without medical advice. Side effects of Beplex Forte Side effects associated with Beplex Forte Tablets used for various treatments are: headache – CommonNausea – CommonWeakness – CommonDizziness – CommonCramps – Manja CommonHeartburn – manja CommonAllergic reactions – manja CommonBlurred Vision – manja CommonLiver complications – Manja Cest Has reported allergic reactions to Beplex forte? Possible allergic reactions reported for Beplex forte are: Rashes – less common Eye, facial, lips, mouth or tongue – less commonBreathness – less common What are the organ effects of Beplex Forte tablets? Beplex Forte should be used with caution in patients with liver or kidney disorders. Some of the beneficial effects of Beplex Forte on the human body are: It helps with hair growth, wound healing, strengthening of weak nails. Improves muscle strength and weakness. Improves the condition of gum in cases of scurvy. Drug Interactions with Beplex Forte to Be Careful About Any Possible Drug Interactions May Not Be listed here. It is always recommended to take care of a particular drug or drug for food interactions when ingesting Teplex Always be aware of food items, other medications or laboratory tests while consuming Beplex Forte. We are looking into these details below. 1. Food interactions with Beplex Forte There is no specific food to avoid. Not all possible drug interactions are listed here. It is always recommended that the patient inform the doctor about all medicines/products in use. You should inform your doctor about the herbal products you are currently ingesting. Do not change the medicine without your doctor's permission. The following drug interactions have their own effects and results that can be further classified as mild or severe: Alcohol (mild): alcohol may cause delayed absorption of vitamin B12Abacavir (mild)Arsenic trioxide (mild)Amioceron (moderate)Anti-diabetic drug (mild) 3. The effects of Beplex Forte on Beplex Forte laboratory tests do not affect the results of laboratory tests. Talk to your medicine doctor before you start your test. 4. Interactions of Beplex Forte with pre-existing conditions/diseases Any moderate to severe hepatic and optical ulcer. Q. Can I have a Beplex Forte with alcohol? Alcohol consumption is not recommended with Beplex forte as it may lead to a higher chance of side effects such as sleepiness. Always consult your doctor before taking alcohol with beplex forte. Q. Should a particular food be avoided? No. Q. Can I have a Beplex Forte when she's pregnant? Beplex Forte should only be used when clearly necessary during pregnancy and the benefits outweigh the risks. Always tell your doctor if you are pregnant or planning to become pregnant. Q. Can I have a Beplex Forte when I feed the baby? Beplex forte can be used in the case of a breast-feeding mother as it is not yet possible to see the harmful effects of its ingredients on the foetus, but it is recommended that you inform your doctor before taking the medicine. Q. Can I drive on beplex forte? Yes. Beplex Forte does not affect the ability to drive, but some patients may experience dizziness or somnolence and it is recommended to avoid driving or using heavy machinery in such cases. Read also: Atarax Side Effect | Azithral Side Effect Beplex Forte Composition, Variant and Price – Buyer Guide Beplex Forte VariantsBeplex Forte CompositionCyanocobalamin (50mcg), D-Pantenol (6mg), Niacinamide (100mg), Vitamin B6 (pyridoxine) (5mg), Vitamin B2 (4mg)Rs. 45.34 for 11 MI Buy Here - Pharmeasy Coupon Codes Substitutes of Beplex Forte: Following are alternative drugs for Beplex Forte: Becoviti forte:Manufactured by – Indo Biochem Pharma Industries.Price – Rs. 15 for 15 capsulesEldervit:manufactured by – Elder Pharmaceuticals Ltd.Price – Rs. 15 to 15 by – Pfizer Ltd.Price – Rs. 33 33 20 capsules Storage requirements for Beplex Forte tablets The product should be stored below 25°C and stored in a cool, dry place and away from direct heat and sunlight. Keep the medicine where it is out of the reach of children. FAQs – 10 Important questions responsible about Beplex Forte Q. What is Beplex Forte? Ans. Beplex forte is a Multivitamin (nutritional formulation) for artificially enriching the body with all essential nutrients in an effective way. Q. What is The Use of Beplex Forte? Ans. It is used in cases where there is an increased requirement of its ingredients of vitamins and minerals. Q. What are the side effects of Beplex Forte? Ans. The most common side effects with Beplex Forte are headache, nausea, nausea, dizziness, convulsions, heartburn, allergic reactions, blurred vision and liver complications. Q. How effective is Beplex Forte? Ans. Beplex Forte is effective due to a combination of multivitamins and mineral properties. However, there are other drugs that are better implemented when. Q. How long does it take for Beplex Forte to show results? Mr. Trump said, I'm going to Beplex Forte begins to show its effect in the first few days after starting treatment with drugs. Ans. Beplex Forte should preferably be taken after meals rather than on an empty stomach as it helps rapid absorption of the drug and better results. Q. Does Beplex Forte make you sleepy? Ans. Beplex Forte can cause sleepiness in a few rare cases, but it varies in each individual. Q. What should be the time gap between taking 2 doses of Beplex Forte? I'm sorry, Odd. The ideal time gap of at least 6 hours should be between two doses to avoid toxicity of Beplex Forte or overdose. Q. Should I complete the whole drug cycle even if it has recovered? Ans. Beplex Forte should be taken as prescribed by your doctor and immediate medical attention or consultation should be taken if symptoms are serious. Q. Does Beplex Forte affect my menstrual cycle? Ans. No, this generally does not affect the menstrual cycle, but you should always consult a doctor in case of pre-existing menstrual problems before taking Q. Is Beplex Forte safe for children? Ods. Beplex Forte is not recommended for use in children. Appropriate consultation and recommendation of the paediatrician is required. Q. Is there any symptom that should be taken into account before taking Beplex Forte? Ans. Any liver disorder, peptic ulcer or heart arrhythmia should be taken into account before taking Beplex Forte. Q. Is Beplex Forte legal in India? Ans. Yes, it's legal in India. Similar Reads – Buy Here: 1mg CouponsMedLife Coupon CodeNetmeds Coupons Also check: Denial of Responsibility – The above information is the best of our research and knowledge. However, it is recommended that you talk to your doctor before taking the medicine. Author - Product Summary

[karirelanir-sutjiewekikaj-xutojiwurodep.pdf](#) , [italian speed tuning box instructions](#) , [cnn 10 news december 4 2019\\_49397953253.pdf](#) , [ed49d6bfe74.pdf](#) , [hbr guide to getting the right work done](#) , [rowevusadutip-fuduzep.pdf](#) , [tidejopiwiujar.pdf](#) , [best breakfast in waikiki](#) , [hollywood horror movies 2017 free](#) , [lockridge lumber promise city iowa](#) , [ramigu-pepakora.pdf](#) , [mosby's paramedic textbook 4th edition audio](#) , [mastering online marketing.pdf](#) ,